

JES-Life Sciences

Protective Effect of Vitamin E on Endosulfan Induced Testicular Toxicity in Swiss Mice (*Mus musculus*)

Girima Nagda*, Habiba Bano and Devendra Kumar Bhatt

Cancer Biology and Toxicology Research Lab, Department of Zoology, M. L. Sukhadia University, Udaipur 313 001, India

Article Info

Article History

Received : 13-01-2011
 Revised : 16-03-2011
 Accepted : 17-03-2011

*Corresponding Author

Tel : +91-9460315237
 Fax : +91-2942425364

Email:
 girima10@gmail.com

Abstract

The present study was aimed to evaluate the role of vitamin E against the testicular toxicity induced by endosulfan. 8-10 weeks old, 24 healthy male animals were randomly selected and equally divided into 4 groups. Group I served as control group (C); group II endosulfan group (ES); group III vitamin E group (VE) and group IV vitamin E plus endosulfan group (VE+ES). Group C animals were given only vehicle the olive oil; in groups ES and VE+ES endosulfan was administered orally at a dose of 2.45 mg/kg b.w.; in groups VE and VE+ES, vitamin E was administered orally at a dose of 50 mg/kg b.w. In group VE+ES, vitamin E was administered 1 hour prior to endosulfan administration. All treatments were given continuously for 15 days. Endosulfan intoxication resulted in decreased testis weight and severe histopathological changes which included shrunken and distorted seminiferous tubules and atrophy in the tissue. It was observed that administration of vitamin E minimized the endosulfan induced damage. Thus it can be concluded that pretreatment with vitamin E can alleviate the damage caused to testis by endosulfan.

©ScholarJournals, SSR

Key Words: Vitamin E; Endosulfan toxicity; Testis; Mice

Introduction

Endosulfan is a potent organochlorine pesticide, under the cyclodiene subgroup that has emerged as a leading chemical used against a broad spectrum of insects and mites in agriculture and allied sectors. India is the world's largest producer and user of endosulfan (1) and it is one of the most commonly used pesticides in India (2). Studies indicate that brain, liver, kidney, immune system and testes are the target organs bearing major impact of endosulfan toxicity (3,4). Endosulfan is known to damage the endocrine system, nervous system, circulatory, reproductive, respiratory and excretory systems and developing foetus (5-8).

Traces of endosulfan have been detected in a great number of supermarket goods including vegetables, seafood and even spices (9). Endosulfan is readily absorbed by humans via the stomach, lungs and through the skin. It also causes endocrine disruption (10).

Endosulfan is known to cause possible damage to reproductive system of male rats (10). In recent years, there has been growing concern about toxicity of pesticides, on the male reproductive system (11,12). It has been reported that endosulfan exposure resulted in delayed sexual maturity in boys in Kerala, India where endosulfan was used on cashew plantations (13).

Vitamin E is one of the most potent lipid soluble free radical scavenger that limits lipid peroxidation initiated by free radicals in the testicular microsomes and mitochondria (14). Vitamin E is a powerful lipophilic antioxidant vital for the maintenance of mammalian spermatogenesis (15). It also reverses the detrimental effects of oxidative stress on testicular

function mediated by different agents (16-19)

Histopathological analysis is regarded as a realistic biomarker in toxicological studies (20). Histopathological studies are useful to evaluate the pollution potential of pesticides since trace levels of pesticides, which do not cause animal mortality over a given period, are capable of producing considerable damage in the tissues (21). Any toxicant either acts directly on the cell or most frequently causes cytotoxicity by modulating the cellular microenvironment. The cells in turn respond histopathologically by degeneration, proliferation, inflammation and repair (22).

In vivo histopathological alterations in the testis due to endosulfan intoxication and the protective role of vitamin E seem to be lacking in the literature. Keeping in mind the above facts, the present study was designed to have an insight into histological analysis regarding the extent of damage of the testicular tissue due to endosulfan toxicity and its subsequent mitigation by Vitamin E.

Materials and Methods

Chemicals

Endosulfan (CAS No. 115-29-7 and purity 99.9%) was obtained from Shree Pesticides India Ltd. Vitamin E was obtained from Himedia, India. All other chemicals and solvents used were of analytical grade.

Animals and Treatment

Male Swiss mice numbering 24, weighing 30 ± 5 gms and 8- 10 weeks old were used in the present study. The animals were maintained on sterilized rice husk bedding in

polypropylene cages and kept at a temperature of about 23 ± 3 °C with 12 ± 1 hour light/dark cycles. The animals were fed on standard pellet diet (Pranav Agro, Baroda). Food and water were given *ad libitum*.

This experimental study was approved by the Institutional Animal Ethics Committee. The handling of the animals was according to the guidelines of Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Ministry of Environment and Forests, Govt. of India.

Dose Selection

The dose for endosulfan was selected after conducting pilot experiments in our laboratory. The LD50 for endosulfan

was found to be 7.35 mg/kg b.w. (23). The dose selected for endosulfan was one third of the LD50 i.e., 2.45 mg/kg b.w. and the duration of treatment was 15 days. The dose selected was lower than which was used in earlier studies (24)

The doses for antioxidants were calculated keeping in mind the doses prescribed for humans. The dose of vitamin E selected was 50 mg/kg b.w.

The doses were prepared by dissolving in olive oil.

Experimental Protocol

A sub chronic study was done for 15 days and oral route of dose administration was chosen for all the treatments. The mice were divided into four groups with minimum of 8-10 animals in each group.

Group I	Control (C) group (given olive oil)
Group II	Endosulfan (ES) group (given 2.45 mg/kg b.w. dose of endosulfan dissolved in olive oil)
Group III	Vitamin E (VE) group (given a vitamin E @ 50 mg/kg b.w.)
Group IV	Vitamin E and endosulfan (VE+ES) group (given vitamin E @ 50 mg/kg b.w plus endosulfan @ 2.45 mg/kg b.w.)

In group IV, the vitamin E was administered 1 hr prior to endosulfan administration.

Histological Preparation

The mice were sacrificed by cervical dislocation at the end of the scheduled period of 60 days and 24 hrs after the last dose treatment. After dissection both the testis were weighed and the left testes was fixed in 10% formalin solution. After 18 to 24 hrs of fixation the testes was further processed for paraffin embedding. Serial sections of 5-7 μ m thick were cut using a rotary microtome. The deparaffinized sections were routinely stained with haematoxylin and eosin, examined and photomicrographed.

Result

Effect on testis weight

A significant reduction in the testis weight was observed in the endosulfan intoxicated animals as compared to the control (C) group. Endosulfan intoxication resulted in a significant decrease ($P < 0.01$) in the testis weight as compared to control (C) group. Pretreatment with vitamin E in group VE+ES significantly alleviated ($P < 0.01$) the testis weight as compared to group ES (Table 1).

Table 1—Changes in the average weight (g) of body weight in control (C), endosulfan treated (ES), Vitamin E (VE) and vitamin E plus endosulfan treated (VE+ES) groups [Values are Mean \pm SD].

Group	Average body weight (mg)
C	119.17 \pm 5.84
ES	97.50 \pm 5.24**a
VE	125.00 \pm 4.47
VE + ES	115.83 \pm 3.76**b

** $P < 0.01$ * $P < 0.05$; a: group ES compared with group C; b: group VE+ES compared with group ES

C: Control group; ES: endosulfan treated group; VE: Vitamin E group; VE+ES: Vitamin E plus Endosulfan treated group

Histopathological findings

Normal architecture was seen in the histological sections of testis of control group (C) (Plate 1, Fig.1). Healthy and normal seminiferous tubules were observed. The seminiferous tubules manifested all the cells of the spermatogenetic series such as spermatogonia (SG), spermatocyte, spermatids, spermatozoa (S), and sertoli cells (SC). The lumen of the tubules was occupied by mature spermatozoa. Healthy blood

vessels (BV) and Leydig cells (LC) were present in the interstitium.

Endosulfan intoxication resulted in the adverse changes in the testicular tissue. Sections of testis from endosulfan group (ES) showed degenerative changes. In few tubules the lumen was completely filled with edematous fluid (Plate 1, Fig 2). Sloughing of the germinal cells lining the seminiferous tubules was a common feature among a number of tubules (Plate 1, fig.3). Majority of the seminiferous tubules were shrunken and had a distorted appearance (Plate 1, Fig. 4).

There was depletion and hypertrophy of germ cells. Lumen was enlarged and binucleated (BN) and trinucleated (TN) cells could be seen in the lumen alongwith dead cells (DC) and spermatogonia.(Plate 1 , Fig, 5; Plate 2, Fig, 6&7)) Oligospermia and complete absence of spermatozoa from the lumen was notable. Plate 2, Fig. 8 shows a giant cell (GC) with fusiform nucleus. Leydig cells and blood vessels were sparse in interstitium.

In group VE (Plate 2, Fig.9) the testis revealed normal structure as observed in group C.

In group VE+ES (Plate 2, Fig. 10) sections were quite similar to those of group C and the pathological changes like giant cells, multinucleated cells, enlarged lumen, distorted seminiferous tubules as observed in group ES were not seen. The structure was more or less restored.




Plate 1: — Histological sections of testis stained by H & E at 400 ×. (Fig. 1)-Section of testis of control group showing normal seminiferous tubules with all the cells of spermatogenetic series. Spermatogonia (SG), Sertoli Cells (SC), Spermatozoa (S), Leydig Cells (LC) and Blood Vessels (BV) can be seen.

Endosulfan treated group [Plate 1, Fig. 2 – 5 and Plate 2, Fig. 6 – 8] (Fig. 2)- shows edematous fluid (EF) in the lumen (L); (Fig. 3)-Dislodging and sloughing of the germ cells; (Fig. 4)- Distorted seminiferous tubules (ST) with loss of germ cells and complete loss of spermatogonia; (Fig. 5)- Dead cells (DC) and germ cells are seen in the lumen;




Plate 2 : (Fig. 6) – Binucleated cell (BN) is seen in the lumen; (Fig. 7)- Shrunken seminiferous tubules showing bi- and tri- nucleated (TN) germ cells; (Fig. 8)- A giant cell (GC) with a fusiform nucleus is seen; (Fig. 9) Section of vitamin E treated group showing normal seminiferous tubules; (Fig. 10)- Section of testis of vitamin E plus Endosulfan treated group showing features close to normal structure.

Discussion

Drastic changes were observed in the histology of testis after endosulfan treatment for 15 days. Occurrence of giant cells (25) and multinucleated giant cells (26-28) in the lumen as observed in the present study have been reported in a number of earlier studies. Such changes may be attributed to failure of cytokinesis and hypertrophy (29). Vacuolations observed are similar to earlier reports (30).

The decreased number of Leydig cells in interstitium in the exposed groups is in accordance with earlier reports on testicular toxicity due to other organochlorine pesticides (29). The inhibition of testicular androgen biosynthesis (31) could be attributed to the present observation.

Oligospermia as observed in the present study is a remarkable observation of testicular toxicity as also reported in earlier studies (29, 31-33)

Dead cells in lumen and the decreased number of germ cells may be due to cytotoxic (34) and apoptotic (24) effect of endosulfan. The reduction and degeneration of germ cells can be a possible reason for the endosulfan induced disruption of the spermatogenic (7) and steroidogenic cycle (31,33). This could also be the reason for delayed sexual maturity as reported in case of Kerala poisoning.

Histopathological changes such as atrophy and decreased number of germ cells may be a reason for decreased testicular weights as observed in the present and earlier reports (29, 35,36) or it can be due to increased protein breakdown due to endosulfan (37).

Such pathological changes can be attributed to oxidative stress induced by endosulfan as reported in earlier studies (6,13, 38, 39). Moreover, overproduction of reactive oxygen species damages vital components of cell, like nucleic acids and proteins which further lead to oligozoospermia and abnormal spermatozoa (29).

Vitamin E supplementation resulted in improvement of the structural alterations in the testis due to endosulfan. This amelioration could be attributed to the capacity of vitamin E to scavenge ROS and reduce the oxidative stress as reported in earlier studies (17, 40)

Thus, on the basis of above findings it can be concluded that pretreatment with vitamin E ameliorates the severe pathological alterations in the testis due to endosulfan exposure. Moreover, it can be said that if occupationally exposed workers supplement their diet with vitamin E the endosulfan induced toxicity could be minimized.

Acknowledgment

The authors are thankful to Shree Pesticides Pvt. Ltd., Udaipur for providing Technical grade Endosulfan as gratis.

References

- [1] Li, Y., and Macdonald, R., 2005. Sources and pathways of selected organochlorine pesticides to the Arctic and the effect of pathway divergence on HCH trends in biota, a review. *Sci Total Environ.* 342, 87-106
- [2] Jayashree, R., and Vasudevan, N., 2007b. Persistence and distribution of endosulfan under field conditions. *Environ Monit Assess.* 131, 475-87.
- [3] Banerjee, B. D., Pal, R., Ahmed, T., Kumar, V., Sanvidhan, G., and Ray, A., 2009. Protective effects of antioxidants against endosulfan induced oxidative stress and immunotoxicity in Albino rats. *Ind. J. Exptl. Biol.* 47, 737-739.
- [4] ATSDR., 2000, Agency of Toxic Substances and Disease Registry. Toxicological Profile for Endosulfan. Atlanta, USA. <http://www.atsdr.cdc.gov/toxprofiles/tp41.html>.
- [5] Paul, V., Balasubrahmaniam, E., Jayakumar, A. R., and Kazi, M., 1995. A sex related difference in the neuro behavioural and hepatic effects following chronic endosulfan treatment in rats. *Eur. J. Pharmacol.* 293, 355-60.
- [6] Naqvi, S. M., and Vaishnavi, C., 1993. Bio Accumulative Potential and toxicity of Endosulfan insecticide to non target animals. *10 5*. 347-61.
- [7] Sinha, N., Narayan, R., Sankar, R., and Saxena, D. K., 1995. Endosulfan induced biochemical changes in testis of rats. *Vet. Human Toxicol.* 37, 547-9.
- [8] Reuber, M.D., 1981. The role of toxicity in the carcinogenicity of Endosulfan. *Sci. Total Environ.* 20, 23-47.
- [9] Strandberg, B., and Hites, R., 2001. Concentration of organochlorine pesticides in wine corks. *Chemosphere.* 44, 729-35.
- [10] Silva, M.H., and Gammon, D., 2009. An assessment of the developmental, reproductive, and neurotoxicity of endosulfan. *Birth Defects Res B Dev Reprod Toxicol.* 86, 1-28.
- [11] Murray, T.J., Lea, R.G., Abramovich, D.R., Haites N.E, Fowler, P.A., 2001. Endocrine disrupting chemicals, effects on human male reproductive health. *Early Pregnancy.* 5, 80-112.
- [12] Sharpe, R.M., 2001. Hormones and testis development and the possible adverse effects of environmental chemicals. *Toxicol Lett.* 120, 221-232.
- [13] Saiyed, H., Dewan, A., Bhatnagar, V., Shenoy, U., Shenoy, R., Rajmohan, H., Patel, K., Kashyap, R., Kulkarni, P., Rajan, B., Lakkad, B., 2003. Effect of endosulfan on male reproductive development. *Environ Health Perspect.* 111(16), 1958-62.
- [14] Gavazza, M. B., and Catala, A., 2006. The effect of alpha-tocopherol on lipid peroxidation of microsomes and mitochondria from rat testis. *Prostaglandins Leukot Essent Fatty Acids.* 74, 247.
- [15] Johnson, F.C., 1979. The antioxidant vitamins. *CRC Crit. Rev. Food Sci. Nutr.* 11, 217-309.
- [16] Verma, R.J., and Nair, A., 2001. Ameliorative effect of vitamin E on aflatoxin induced lipid peroxidation in the testis of mice, *Asian J Androl.* 3, 217.
- [17] Latchoumycandane, C., and Mathur, P. P., 2002. Effects of vitamin E on reactive oxygen species- mediated 2,3,7,8-tetrachlorodi-benzo-p-dioxin toxicity in rat testis. *J Appl Toxicol.* 22, 345.
- [18] Gupta, R., Sen Gupta, E., Dhakal, B.K., Thakur, A.R & Ahnn, J., 2004. Vitamin C and vitamin E protect the rat testes from cadmium-induced reactive oxygen species, *Mol Cells.* 17, 132.
- [19] Jedlinska-Krakowska M., and Bomba G *et al.*, 2006. Impact of oxidative stress and supplementation with vitamins E and C on testis morphology in rats, *J Reprod Dev.* 52. 203.
- [20] McCarthy. J.F., and Shugart, L.R., 1990. Biomarkers of environmental contamination. Lewis Pubs., Boca Raton, Florida, USA.
- [21] Kumar, S. and Pant, S.C., 1984. Organal damage caused by aldicarb to freshwater teleost, *Barbus conchonioides* Hamilton. *Bull. Environ. Contam. Toxicol.* 33, 50-53.
- [22] M.S. Prashanth., 2011. Histopathological Changes Observed In The Kidney Offreshwater Fish, *Cirrhinus Mrigala* (Hamilton) Exposed To Cypermethrin. *Recent Research in Science and Technology.* 3(2), 59-6.
- [23] Gupta, P.K., Murthy, R.C., and Chandra, S.V., 1981. Toxicity of endosulfan and manganese chloride: Cumulative toxicity rating. *Toxicol Lett.* 7:221-227.
- [24] Singh, J.K., Ali Mohammad, Kumar Ranjit, Nath, A., Kumar, Arun., 2010. Possible occurrence of apoptosis in Endosulfan induced testicular toxicity in mice" *Journal of Ecophysiology and Occupational Health Year*, 10.
- [25] Chinoy N J., Sorathia, H. P., Jhala, D .D., 2005. Fluoride+Aluminium induced toxicity in mice testis with giant cells and its reversal by vitamin C. *Fluoride.* 38, 109.
- [26] Chowdhury, A. R., Venkatakrishna-Bhatt, H., and Gautam, A. K., 1987. Testicular changes of rats under lindane treatment. *Bull Environ Contam Toxicol.* 38, 154.
- [27] Dikshith, T. S. S., and Datta, K. K., 1972b. Pathologic changes induced by pesticides in the testes and liver of rats. *Exp Path.* 7, 309.
- [28] Nigam, S. K., Lakkad, B. C., Karnik, A. B., Thakore, K. N., Bhatt D K, Arvinda Babu , K., and Kashyap, S. K., 1979.

- Effect of hexachlorocyclohexane feeding on testicular tissues of pure inbred Swiss mice, Bull Environ Contam Toxicol. 23. 431
- [29] Nagda, G., and Bhatt, D.K., 2011. Alleviation of lindane induced toxicity in testis of Swiss mice (*Mus musculus*) by combined treatment with vitamin C, vitamin E and α -Lipoic acid. Ind. J. Exp. Bio. 49, 191-199.
- [30] Eweka, A. O., and Om'Iniabohs, F.A.E., 2008. Histological studies of the effects of monosodium glutamate on the testis of adult Wistar rats. The Internet Journal of Urology. 5.2.
- [31] Singh S.K. and Pandey R.S., 1990. Effect of subchronic Endosulfan exposure on plasma gonadotrophins, testosterone, testicular testosterone and enzyme of androgen biosynthesis in rat. Ind. J. Exp. Biol, 28, 953-956.
- [32] Chitra, K.C., Latchoumycandane, C., Mathur, P.P, 1999. Chronic effect of endosulfan on the testicular functions of rat *Asian J Androl.* 1, 203-206.
- [33] Singh ,S.K., and Pandey, R.S., 1989. Toxicity of endosulfan on kidney of male rats in relation to drug metabolizing enzymes and microsomal lipid peroxidation. Ind. J. Exp. Biol. 27, 725-728.
- [34] Shimoga Durgojirao Manjula., Susan Benjamin and Kurady Laxminarayana Bairy., 2006. Modulatory Effect of Vitamin C on Genotoxic Effect of Endosulfan in Developing Albino Rats *Iranian Journal of Pharmacology & Therapeutics*, 5, 113-116
- [35] Brooks DE., 1983. Effect of androgen on protein synthesis and secretion in various regions of the rat epididymis, as analysed by two-dimensional gel electrophoresis. Mol Cell Endocrinol. 29, 255-70.
- [36] Sujatha, R., Chitra, K.C., Latchoumycandane, C., and Mathur, PP., 2001. Effect of Lindane on testicular antioxidant system and steroidogenic enzymes in adult rats. Asian J Androl. 3, 135.
- [37] Arshad, N., Shabbir, G., Aleem, S., and Arshad, M., 2007. Effects of α -Tocopherol on Liver Biochemistry of Endosulfan Intoxicated Mice, A Preliminary Study. Asian J. Exp. Sci. 21, 239-246.
- [38] Dorval, J., Hontela, A., 2003. Role of glutathione redox cycle and catalase in defense against oxidative stress induced by endosulfan in adrenocortical cells of rainbow trout (*Oncorhynchus mykiss*). Toxicol Appld Pharmacol. 192(2),191-200.
- [39] Bebe, F.N. and M. Panemangalore., 2003. Exposure to low doses of endosulfan and chlorpyrifos modifies endogenous antioxidants of rats. J. Environ. Sci. Health B. 38, 349-363.
- [40] Kelle, M., H. Diken, A. Sermet, M. Atmaca and C. Tumer., 1999. Effect of exercise on blood antioxidant status and erythrocyte lipid peroxidation, Role of dietary supplementation of vitamin E. Turk. J. Med. Sci. 29, 95-100.